



YaraVita™ Bud Builder™ WP

A formulated powder mixture of magnesium and phosphorus with zinc and boron

Guaranteed Analysis:

Available Phosphate (P ₂ O ₅)	6%
Magnesium (Mg)	24%
Boron (B)	3%
Zinc (Zn)	10%

Derived from Zinc Phosphate, Magnesium Hydroxide, Sodium Borate

Benefits:

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com/yara for details
- Proven, reliable performance. Trialled and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- High nutrient content means lower application rates reducing handling time and waste packaging



Product Recommendations

Typical Crop Recommendations*

- **Apples, Pears:** 10 lbs/acre applied post-harvest before leaf fall. Also, 5 to 10 lbs/acre applied at bud burst. Water rate: 50 to 100 gallons/acre.
- **Apricots, Cherry, Nectarines, Peaches, Plums:** 10 lbs/acre applied post-harvest before leaf fall. Repeat if necessary leaving 10 to 14 days between applications. Also, 5 to 10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.
- **Blueberries:** Apply 5 lbs/acre just before onset of leaf drop. Repeat the application at bud separation the following fruiting season. Water rate: 20 gallons/acre.
- **Caneberries:** 5 lbs/acre applied from green bud to just prior to start of flowering. Also, 5 lbs/acre post-harvest, pre-leaf senescence. Water rate: 20 to 50 gallons/acre.
- **Conifers:** 2 applications of 5 to 10 pounds/acre once there is new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.
- **Grapevines:** 1 to 2 applications of 5 lbs/acre applied pre-flowering at a 10-14 day interval. Also, 10 lbs/acre applied post-harvest before leaf fall. Water rate: 50 to 100 gallons/acre.
- **Nuts (Deciduous):** 10 lbs/acre applied post-harvest before leaf fall. Also, 5 to 10 lbs/acre applied from bud burst to start of flowering. Water rate: 50 to 100 gallons/acre.
- **Strawberries (Field Grown):** 5 lbs/acre applied from green bud to just prior to start of flowering. Also, 5 lbs/acre post-harvest, pre-leaf senescence. Water rate: 20 to 50 gallons/acre.

*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.



Yara North America, Inc.
1-800-234-9376
www.yara.us