

Nutra Spray Manzinc

Wettable powder foliar nutrition product.

Guaranteed Analysis:

Cocionico de l'incignio	
Zinc (Zn)	25%
Manganese (Mn)	25%

Benefits:

- Wettable powder that provides metered release of nutrients over several days giving a cost effective treatment.
- Extremely fine particle size gives thorough coverage, efficient nutrient uptake, and enhanced sprayability.
- Offers increased crop safety to even the most sensitive crops. Safe to foliage and young fruit at recommended rates.
- Tank-mixable with most commonly used agrichemicals.







*The information provided is accurate to the best of Yara's knowledge and belief.
Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.



Product Recommendations

Typical Crop Recommendations*

- Alfalfa: Apply to new growth in spring and after each cutting at the rate of 2 to 5 lb per acre per application. Repeat as needed.
- Deciduous Nuts (Almond, Filbert/Kentish Cob Nuts, Pecans, Pistachios, Walnuts): 10 to 30 lb per acre for mature trees. In horticultural oil sprays, 3/4 to 1 lb. may be included for each 1% of oil emulsion.
- Apple, Pears: 10 to 30 lb per acre per application at pre-bloom and post-harvest.
 Apply foliar sprays as needed. In horticultural oil sprays, 3/4 to 1 lb. may be included for each 1% of oil emulsion.
- Stone Fruit (Apricot, Cherry, Nectarines, Peach, Plum, Prunes): 10 to 30 lb per acre for mature trees. In horticultural oil sprays, 3/4 to 1 lb. may be included for each 1% of oil emulsion.
- **Bananas:** 10 to 30 lb per acre per application. Repeat as needed.
- Beans, Peas, Soya Bean: Apply at seedling to pre-bloom stage at a rate of 2 to 4 lb. per acre. Repeat as needed.
- Blueberries, Raspberry: Use 2 to 4 lb. per acre. Begin application before bloom. Repeat as needed. Application after fruit starts to color may leave a visible residue.
- Citrus: 16 to 32 lb. per acre for mature trees.
 Application may be made at any time of year, although for greater efficiency, application should be made over new growth flushes. In horticultural oil sprays, use 1 to 1-1/2 lb. per 100 gal. of finished spray.
- Coconut: 10 to 30 lb. per acre per application. Repeat as needed.
- Cotton: 2 to 10 lb. per acre. First application applied prior to main bloom. Repeat as
- Curcurbits (Field Grown) Courgette,
 Cucumber, Melon, Water Melon: Apply at seedling to pre-bloom stage at the rate of 2 to 4 lb. per acre in enough water to give even coverage.
- Durian: 10 to 30 lb. per acre per application.
 Repeat as needed.
- Grapevines: Apply 2 to 10 lb. per acre prior to bloom. Do not apply later than buckshot stage.
- Guava: 10 to 30 lb. per acre per application.
 Repeat as needed.
- Lentil: Apply at seedling to pre-bloom stage at a rate of 2 to 4 lb. per acre. Repeat as needed.
- Lettuce (Field Grown): 3 to 5 lb. per acre per application. Repeat as needed.
- Longan: 10 to 30 lb. per acre per application.
 Repeat as needed.

- **Lychee:** 10 to 30 lb. per acre per application. Repeat as needed.
- Macadamia: 10 to 30 lb. per acre per application, pre-bloom and in cover sprays. Repeat as needed.
- Maize: 2 to 4 lb. per acre. Apply at any period between the seedling stage and tasseling. Repeat as needed.
- Mango: 10 to 30 lb. per acre per application. Repeat as needed.
- Mangosteen: 10 to 30 lb. per acre per application. Repeat as needed.
- Papaya/Paw Paw: 10 to 30 lb. per acre per application. Repeat as needed.
- Pepper (Field Grown): 2 to 4 lb. per acre at the seedling to pre-bloom stage.
- Pineapple: 10 to 30 lb. per acre per application. Repeat as needed.
- Rambutan: 10 to 30 lb. per acre per application. Repeat as needed.
- Star-fruit: 10 to 30 lb. per acre per application. Repeat as needed.
- Sugar Apple: 10 to 30 lb. per acre per application. Repeat as needed.
- Sugar Beet: 2 to 4 lb. per acre per application. Repeat as needed.
- Tomato (Field Grown): 2 to 4 lb. per acre at the seedling to pre-bloom stage.
- Aerial Applications: Apply aerial applications in at least 5 gallons/acre and ground applications in at least 10 gallons/acre; however, when total application gallonage is 25 gallons per acre or less, DO NOT mix more than 1 lb. of this product per gallon of water.

*The information provided is accurate to the best of Yara's knowledge and belief.

and belief.

Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.

Yara North America, Inc. www.yara.us Tel: 1 800 234 9376

