

Nutra Spray Zn Plus

Wettable powder foliar nutrition product.

Guaranteed Analysis:	
Zinc (Zn)	50%
Calcium (Ca)	7.4%
Sulfur (S)	6%

Benefits:

- High analysis, pH neutral, wettable powder that provides metered release of nutrients over several days giving a cost effective treatment.
- Extremely fine particle size gives thorough coverage, efficient nutrient uptake, and enhanced sprayability.
- Offers increased crop safety to even the most sensitive crops. Safe to foliage and young fruit at recommended rates.
- Tank-mixable with most commonly used agrichemicals and Leffingwell Sorba-Spray products.
- High quality consistent product. Manufactured to ISO 9001 assurance standards.





^{*}The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.



Product Recommendations

Typical Crop Recommendations*

- Apricot, Almond, Cherry, Nectarine, Peach, Plum, Prune: Dormant: Use 6 to 15 lb per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb per acre. May be used with horticultural oil. Foliage Spray: Apply at the rate of 2 to 6 lb per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb per acre.
- Apple: Dormant: Use 6 to 15 lb per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb per acre. May be used with horticultural oil.
 Foliage Spray: Apply at the rate of 2 to 6 lb per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb per acre.
- **Avocado:** 5 to 15 lb per acre per application. Repeat as needed.
- Bananas: 2 to 6 lb per acre per application. Repeat as needed.
- Beans: 2 to 5 lb. per acre per application. Repeat as needed.
- **Cereals:** 2 to 5 lb. per acre per application. Repeat as needed.
- Citrus: 10 to 20 lb. per acre per application for mature trees. In horticultural oil sprays, use 3/4 to 1 lb. per 100 gal. in dilute sprays; 5 to 10 lb. per acre in concentrate sprays.
- **Coconut:** 2 to 6 lb. per acre per application. Repeat as needed.
- Durian: 2 to 6 lb. per acre per application. Repeat as needed.
- Filbert/Kentish Cob Nuts: Dormant: Use 6 to 15 lb. per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb. per acre. May be used with horticultural oil. Foliage Spray: Apply at the rate of 2 to 6 lb. per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb. per acre.
- **Grapevines:** Apply 2 to 8 lb. per acre applied at 4-6" new shoot growth, not later than buckshot stage. May be used in a dormant spray at the rate of 3/4 lb. per gal. of horticultural oil.
- Guava: 2 to 6 lb. per acre per application. Repeat as needed.
- Lentil: 2 to 5 lb. per acre per application. Repeat as needed.
- Longan: 2 to 6 lb. per acre per application. Repeat as needed.

- Lychee: 2 to 6 lb. per acre per application. Repeat as needed.
- Maize: 3 to 5 lb. per acre per application. Repeat as needed.
- Mango: 2 to 6 lb. per acre per application. Repeat as needed.
- **Mangosteen:** 2 to 6 lb. per acre per application. Repeat as needed.
- Millet: 2 to 5 lb. per acre per application. Repeat as needed.
- Papaya / Paw Paw: 2 to 6 lb. per acre per application. Repeat as needed.
- Pears: Dormant: Use 6 to 15 lb. per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb. per acre. May be used with horticultural oil. Foliage Spray: Apply at the rate of 2 to 6 lb. per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb. per acre. Apply aerial applications in at least 5 gallons/acre; however, when total application gallonage is 25 gallons per acre or less, DO NOT mix more than 1 lb. of this product per gallon of water.
- Peas: 2 to 5 lb. per acre per application. Repeat as needed.
- Pecans: Dormant: Use 6 to 15 lb. per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb. per acre. May be used with horticultural oil. Foliage Spray: Apply at the rate of 2 to 6 lb. per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb. per acre.
- **Pineapple:** 2 to 6 lb. per acre per application. Repeat as needed.
- Pistachios: Dormant: Use 6 to 15 lb. per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb. per acre. May be used with horticultural oil. Foliage Spray: Apply at the rate of 2 to 6 lb. per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb. per acre.
- **Potatoes:** 2 to 5 lb. per acre per application.
- **Rambutan:** 2 to 6 lb. per acre per application. Repeat as needed.
- Rice: 2 to 5 lb. per acre per application. Repeat as needed.

- **Sorghum:** 2 to 5 lb. per acre per application. Repeat as needed.
- Star-fruit: 2 to 6 lb. per acre per application. Repeat as needed.
- **Sugar Apple:** 2 to 6 lb. per acre per application. Repeat as needed.
- Walnuts: Dormant: Use 6 to 15 lb. per acre. May be used with horticultural oil. Delayed Dormant: Use 5 to 10 lb. per acre. May be used with horticultural oil. Foliage Spray: Apply at the rate of 2 to 6 lb. per acre per application. Repeat as needed. Application of this product as a foliage spray must be made with caution, especially during wet weather or poor drying conditions. Application to fruit within 3 weeks of harvest may leave a visible residue. Post Harvest: 6 to 15 lb. per acre.

*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.

> Yara North America, Inc. www.yara.us Tel: 1 800 234 9376

