



B-Moly[™]



NOTICE

Before using this product, read the entire label

B-Moly™

3-2-0

A concentrated liquid fertilizer with micronutrients.

Guaranteed analysis:	
Total Nitrogen (N)	3%
3% Urea Nitrogen	
Available Phosphate (P2O5)	2%
Boron (B)	

Derived from urea, phosphoric acid, boric acid, sodium molybdate

Density: 10.7 lbs/gal @ 68º F

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

Danger

H360: May damage fertility or the unborn child. P280: Wear protective gloves/clothing and eye/face protection. P202: Do not handle until all safety precautions have been read and understood. P308: IF exposed or concerned: P313: Get medical attention. P501: Dispose of contents and container according to local regulations.



F1106

Guaranteed by **Yara North America, Inc.** 100 North Tampa Street, Suite 3800, Tampa, FL 33602 Customer Service: 800-234-9376

Contents made in Great Britain.

Manufactured by **Yara UK Ltd.**, Manor Place, Pocklington, York, YO42 1DN. U.K.

ONLY APPLY TO THE INTENDED APPLICATION SITE. DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES OR TO ANY IMPERVIOUS SURFACE. DO NOT APPLY IF HEAVY RAIN IS EXPECTED WARNING - This product contains Boron (B) and Molybdenum (Mo) which may be injurious to certain crops and toxic to ruminant animals. Contact your county agent or field consultant for specific information.

APPLICATION RATES AND TIMINGS

Cotton: 1-2 pints/acre at 4 to 6 leaf stage, at appearance of first flower bud squares and again at open flowers stage. Water rate: 5 to 15 gallons/acre.

Maize, Corn: 1-2 pints/acre at 4 to 8 leaf stage. For moderate to severe deficiency, a repeat application may be necessary 10 to 14 days later. Water rate: 10 to 20 gallons/acre.

Potatoes: One to two applications of 1-2 pints/acre applied from 7 to 14 days after 100 percent emergence to 20 days after tuber initiation and, following petiole analysis, during tuber bulking.

Water rate: 10 to 20 gallons per acre.

Soybeans: 1-2 pints/acre when crop is 2 to 6 inches tall, repeated at 10 to 14 day intervals if necessary. Water rate: 5 to 20 gallons/acre.



Extended recommendations available at www.yara.us/crop-nutrition/fertilizer-products/yaravita

DIRECTIONS FOR USE.

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures ≤ 23°F and > 95°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

 \triangle

WLUSYPCTL01