

## **YaraVita**<sup>™</sup>

Bud Builder™WP



#### NOTICE

Before using this product, read the entire label

# Bud Builder™WP

A formulated powder mixture of magnesium and phosphorus with boron and zinc



THIS PRODUCT IN NON-BULK PACKAGING IS A MARINE POLLUTANT WHEN TRANSPORTED BY VESSEL. HOWEVER, IT IS NOT DOT REGULATED WHEN TRANSPORTED BY HIGHWAY ONLY PER 49 CFR, 171.4(c)

#### Danger

H360: May damage fertility or the unborn child. H411: Toxic to aquatic life with long lasting effects. P202: Do not handle until all safety precautions have been read and understood. P280: Wear protective gloves/clothing and seye/face protection. P273: Avoid release to the environment. P308+P313: IF coexposed or concerned: Get medical attention. P391: Collect spillage. P501: Dispose of contents and container according to local regulations.

#### 1106

Guaranteed by Yara North America, Inc.

100 North Tampa Street, Suite 3800, Tampa, FL 33602

Customer Service: 800-234-9376 Contents made in Great Britain.

Manufactured by Yara UK Ltd., Manor Place, Pocklington, York, YO42 1DN, U.K.

### YaraVita™Bud Builder™WP

#### WARNING

This product contains boron which may be injurious to certain crops. This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis. Contact your county agent or field consultant for specific information.

ONLY APPLY TO THE INTENDED APPLICATION SITE. DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES OR TO ANY IMPERVIOUS SURFACE. DO NOT APPLY IF HEAVY RAIN IS EXPECTED.

#### APPLICATION RATES AND TIMINGS

Apples, Pears\*: In the Spring apply 5 to 10 lbs/acre applied at bud burst. Reapply after fruit set during sizing in 10 to 14 day intervals. 5-10 lbs/acre applied post-harvest before leaf senescence. Water rate: minimum 25 gallons/acre.

Apricots, Nectarines Peaches, Cherries, Prunes\*: 5-10 lbs/acre. Apply from bud burst to start of flowering. Apply petal fall thru early fruit sizing in 10-14 day intervals thru green fruit sizing & leaf canopy development period. Apply post-harvest in 21-28 day interval till leaf senescence. Water rate: minimum 25 gallons/acre.

Blackberry, Blueberry, Boysenberry\*: Apply 3-5 lbs/acre: Apply from bud swell thru pink bud. Apply post bloom-petal fall thru green berry sizing and leaf development period in 14 day intervals. Reapply post-harvest during fruit bud formation, in 21-28 day interval until the onset of leaf senescence. Water rate: minimum 20 gallons/acre.

Brassicas\*: 5-10 lbs per acre at the 4 to 6 leaf stage, with repeat applications at 10 to 14 day intervals.

Water rate: minimum 20 gallons per acre.

Citrus, Oranges\*: 5-10 lbs/acre applied post-harvest. Also, 5-10 lbs/acre applied from bud burst to start of flowering Reapply from petal fall thru fruit sizing in 14-21 day interval.

Water rate: minimum 25 gallons per acre.

**Conifers:** 2 applications of 5-10 pounds/acre once there is new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.

Grapevines\*: Apply 1 to 3 applications of 5-10 lbs/acre applied bud break thru pre-bloom at a 10-14 day interval. Also, during post-bloom thru berry sizing 1-2 applications in 7-14 day intervals. Also apply post-harvest before leaf senescence. Water rate: minimum 25 gallons/acre.

Nuts (Deciduous), Almonds, Hazelnuts, Pistachios, Pecans, Walnuts: Apply 5 to 10 lbs/ac applied from bud burst to start of flowering, After nut set, repeat in 14-28 day intervals. Apply 5-10 lbs/ac post-harvest before leaf senescence. Water rate: 25 gal/ac minimum.

Potatoes: 5 lbs/acre one week after plant emergence. Repeat at 10 to 14 day intervals. Water rate: 20 gallons per acre.

Raspberry, Strawberry: 5 lbs/acre applied from green tip to pink bud to just prior to start of flowering in 10-14 day intervals. Apply 1 to 3 times from petal fall thru berry sizing. Also, 5 lbs/acre post-harvest in 21 to 28 day interval up to leaf senescence. Water rate: minimum 20 gallons/acre.

\*Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.



Extended recommendations available at www.yara.us/crop-nutrition/fertilizer-products/yaravita

#### DIRECTIONS FOR USE.

#### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

#### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

#### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com

#### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures ≤ 23°F and > 95°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

#### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request.

Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

