



# YaraVita™

## Bud Builder™ WP 0-6-0

### NOTICE

Before using this product, read the entire label

**A formulated powder mixture of magnesium and phosphorus with zinc and boron**

Guaranteed analysis:

Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....	6%
Magnesium (Mg).....	24%
0.00% water soluble Magnesium (Mg)	
Boron (B).....	3%
Zinc (Zn).....	10%

Derived from Zinc Phosphate, Magnesium Hydroxide, Sodium Borate

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd., Manor Place, Wellington Road,  
The Industrial Estate, Pocklington, York YO42 1DN, U.K.  
Manufactured for Yara North America, Inc. 100 North Tampa Street, Suite 3200, Tampa,  
FL 33602. Customer Service: 800-234-9376

Information regarding the contents and levels of metals in this product is available on the  
internet at <http://www.aapfco.org/metals.htm>

This product is recommended for use in conjunction with a complete fertilization program  
based on soil or tissue analysis.

**Net weight 22lb / 9.9 kg**

F1673

**THIS PRODUCT IN NON-BULK PACKAGING IS A MARINE POLLUTANT WHEN TRANSPORTED BY VESSEL.  
HOWEVER, IT IS NOT DOT REGULATED WHEN TRANSPORTED BY HIGHWAY ONLY PER 49 CFR, 171.4(c)**



UN 3077

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (Irrizinc bis(orthophosphate))

### Danger

Contains disodium octaborate

H360: May damage fertility. May damage the unborn child.

H411: Toxic to aquatic life with long lasting effects.

P201: Obtain special instructions before use.

P202: Do not handle until all safety precautions have been read and understood.

P280: Wear protective gloves/clothing and eye/face protection.

P308+P313: IF exposed or concerned: Get medical attention.

P501: Collect spillage.


P501: Dispose of contents and container according to local regulations.



5H4/Z11/S/16/GB/6550

10.0617

WAL09981



# YaraVita™ Bud Builder WP™

## WARNING

This fertilizer contains boron which may be injurious to certain crops. Contact your county agent or field consultant for specific information.

## APPLICATION RATES AND TIMINGS

**Apples, Pears:** In the Spring apply 5 to 10 lbs/acre applied at bud burst. Reapply after fruit set during sizing in 10 to 14 day intervals. 5-10 lbs/acre applied post-harvest before leaf senescence. Water rate: minimum 25 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

**Apricots, Nectarines Peaches, Cherries, Prunes:** 5-10 lbs/acre. Apply from bud burst to start of flowering. Apply petal fall thru early fruit sizing in 10-14 day intervals thru green fruit sizing & leaf canopy development period. Apply post-harvest in 21-28 day interval till leaf senescence. Water rate: minimum 25 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

**Blackberry, Blueberry, Boysenberry:** Apply 3-5 lbs/acre: Apply from bud swell thru pink bud. Apply post bloom-petal fall thru green berry sizing and leaf development period in 14 day intervals.

Reapply post-harvest during fruit bud formation, in 21-28 day interval until the onset of leaf senescence. Water rate: minimum 20 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

**Brassicas:** 5-10 lbs per acre at the 4 to 6 leaf stage, with repeat applications at 10 to 14 day intervals. Water rate: minimum 20 gallons per acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

**Citrus, Oranges:** 5-10 lbs/acre applied post-harvest. Also, 5-10 lbs/acre applied from bud burst to start of flowering Reapply from petal fall thru fruit sizing in 14-21 day interval. Water rate: minimum 25 gallons per acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

**Conifers:** 2 applications of 5-10 pounds/acre once there is new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.

**Grapevines:** Apply 1 to 3 applications of 5-10 lbs/acre applied bud break thru pre-bloom at a 10-14 day interval. Also, during post-bloom thru berry sizing 1-2 applications in 7-14 day intervals. Also apply post-harvest before leaf senescence. Water rate: minimum 25 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

**Nuts (Deciduous), Almonds, Hazelnuts, Pistachios, Pecans, Walnuts:** Apply 5 to 10 lbs/ac applied from bud burst to start of flowering. After nut set, repeat in 14-28 day intervals. Apply 5-10 lbs/ac post-harvest before leaf senescence. Water rate: 25 gal/ac minimum.

**Potatoes:** 5 lbs/acre one week after plant emergence. Repeat at 10 to 14 day intervals. Water rate: 20 gallons per acre.

**Raspberry, Strawberry:** 5 lbs/acre applied from green tip to pink bud to just prior to start of flowering in 10-14 day intervals. Apply 1 to 3 times from petal fall thru berry sizing. Also, 5 lbs/acre post-harvest in 21 to 28 day interval up to leaf senescence. Water rate: minimum 20 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

*Extended recommendations available at*  
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

## DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

## CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



**WARNING: Cancer and reproductive Harm - [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)**