



Knowledge grows

YaraVita[®] COPTRAC[™]

A formulated product for the treatment of copper deficiency by foliar or soil application

Guaranteed Analysis	
Total Nitrogen (N)	4%
Urea Nitrogen	4%
Copper (Cu)	23.8%
Derived from Urea, Copper Oxide	

The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions.



Benefits

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com/yara for details.
- Proven, reliable performance. Tried and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank.
- High nutrient content means lower application rates reducing handling time and waste packaging

Product Recommendations

Typical Crop Recommendations*

- **Alfalfa:** 0.5 pint/acre applied at start of tillering. Water rate: 20 gallons/acre
- **Carrots:** ½ pint/acre when crop is 6" tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 20 gallons/acre.
- **Cereals:** ½ pint/acre from 2 leaf stage to second node detectable (Zadok's G.S. 12 to 32). For fall-planted crops, an application prior to winter dormancy is strongly recommended. For moderate to severe deficiency, repeat the application once or twice at 10-14 day intervals. Water rate: 20 gallons/acre.
- **Conifers:** 2 applications of ½ pint/acre once there is new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.
- **Corn:** ½ pint/acre at 4 to 6 leaf stage. Water rate: 20 gallons/acre.
- **Garlic:** ½ pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6 inches tall. Repeat applications at 10 to 14 day intervals. Water rate: 7.5 to 20 gallons/acre.
- **Turf:** 0.2 fl.oz./1000 sq.ft. as soon as growth commences in spring. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 0.5 gallons/1000 sq.ft.
- **Nursery/Ornamentals:** One pint in 100 gallons water (0.125% v/v) as soon as there is sufficient leaf area to intercept a spray. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Spray a maximum of three applications per crop per annum. Note: Do not apply within one month of picking/marketing. Maximum water rate: 20 gallons/acre.
- **Onions:** ½ pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6" tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 20 gallons/acre.
- **Potatoes:** 0.5 pint/acre applied 7 to 14 days after 100% emergence. Water rate: 20 gallons/acre
- **Rice:** ½ pint/acre applied at start of tillering. Water rate: 20 gallons/acre.
- **All Crops Soil Application:** 2 quarts/acre to 1 gallon/acre, applied pre-planting or pre-emergence. Water rate: 5 gallons/acre minimum
- **Sorghum:** ½ pint/acre at the 4 to 8 leaf stage. Water rate: 3 to 20 gallons/acre.
- **Sweet Corn:** ½ pint/acre at 4-6 leaf stage. Water rate: 20 gallons/acre.

*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.