



Knowledge grows

# YaraVita<sup>®</sup> MAGPHOS<sup>®</sup>

A formulated product for the treatment of magnesium and phosphorus deficiencies by foliar application

## Guaranteed Analysis

Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	29%
Soluble Potash (K <sub>2</sub> O)	5%
Magnesium (Mg)	4.1%
Water soluble Magnesium (Mg)	4.1%

Derived from Phosphoric Acid, Magnesium Phosphate, Potassium Chloride, Potassium Phosphate

The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions.



## Benefits

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit [www.tankmix.com/yara](http://www.tankmix.com/yara) for details.
- Proven, reliable performance. Tried and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank
- High nutrient content means lower application rates reducing handling time and waste packaging

# Product Recommendations

## Typical Crop Recommendations\*

- **Alfalfa:** 2 quarts/acre when sufficient leaf cover to intercept spray. Then 2 quarts/acre one week after every cut. Water rate: 20 gallons/acre.
- **Apples, Pears:** 4 to 6 applications of 2 quarts/acre from end of flowering. Repeat applications at 10 to 14 day intervals. Water rate: 50 to 100 gallons/acre.
- **Apricots:** 2 to 5 applications of 1½ to 2 quarts/acre from petal fall. Repeat applications at 10 to 14 day intervals. Water rate: 100 gallons/acre.
- **Blueberries:** 2 quarts/acre 10 days after petal fall has finished. Water rate: 100 gallons/acre Brassicas,
- **Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower:** 2 quarts/acre applied soon after transplanting. Water rate: 20 gallons/acre.
- **Carrots:** 2 quarts/acre when crop is 6 inches tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Water rate: 20 gallons/acre
- **Cereals:** 2 quarts/acre at tillering. Repeat at 10 to 14 day intervals if necessary. Also apply from ear emergence until the end of flowering (Zadok's G.S. 51 to 69). Water rate: 20 gallons/acre.
- **Cherries:** 2 to 5 applications of 1½ to 2 quarts/acre from petal fall. Repeat applications at 10 to 14 day intervals. Water rate: 100 gallons/acre.
- **Corn:** 2 quarts/acre at 4 to 8 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 3 to 20 gallons/acre.
- **Cotton:** 2 quarts/acre at the start of flowering. Water rate: 5 to 20 gallons/acre.
- **Cucurbits (Field Grown):** 2 quarts/acre at 4 leaf stage. Repeat if necessary 10 to 14 days later. Water rate: 50 gallons/acre.
- **Ginseng:** 2 quarts/acre applied once the new season growth is well underway. Repeat applications at 10 to 14 day intervals may be necessary for moderate to severe deficiency. Water rate: 50 gallons/acre
- **Vines:** 1½ to 2 quarts/acre at flower buds separated, fruit set, grape expansion and early veraison/one month before harvest. Water rate: 20 gallons/acre.
- **Turf:** 1.5 fl.oz./1000 sq.ft. as soon as growth commences in spring. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 0.5 gallons/1000 sq.ft.
- **Groundnuts:** 2 quarts/acre at the 4 to 6 leaf stage. Repeat as necessary at 10 to 14 day intervals. Water rate: 20 gallons/acre.
- **Lettuce (Field Grown):** 2 quarts/acre. Apply 10 to 14 days after transplanting or emergence. One or two repeat applications may be made at 10 to 14 day intervals. Water rate: 50 gallons/acre
- **Melons (Field Grown):** 1½ to 2 quarts/acre at start of flowering. Repeat at 10 to 14 day intervals if necessary. Water rate: 50 gallons/acre
- **Nectarines:** 2 to 5 applications of 1½ to 2 quarts/acre from petal fall. Repeat applications at 7 to 10 day intervals. Water rate: 100 gallons/acre.
- **Nursery/Ornamentals:** 2 gallons in 100 gallons water (1.875% v/v) as soon as there is sufficient leaf area to intercept a spray. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Spray a maximum of three applications per crop per annum. Maximum water rate: 20 gallons/acre
- **Canola:** For a single application, 1½ to 2 quarts/acre at onset of stem extension. For moderate deficiency, 1½ to 2 quarts/acre at 4 to 6 leaf stage and at onset of stem extension. An extra application can be made 10 to 14 days later for severe deficiency. Avoid flowering. Water rate: 20 gallons/acre.
- **Onions:** 2 quarts/acre. Apply when the foliage is 6" tall, with a second application 10 to 14 days later if necessary. Water rate: 20 gallons/acre
- **Peaches:** 2 to 5 applications of 1½ to 2 quarts/acre from petal fall. Repeat applications at 10 to 14 day intervals. Water rate: 100 gallons/acre.
- **Peas:** 1 to 2 applications of 2 quarts/acre when the crop is 4 to 6" tall. Allow 10 to 14 days between applications. Water rate: 20 gallons/acre.
- **Peppers (Field Grown):** 1 pint in 45 gallons before transplanting. Water rate: 100 gallons/acre maximum
- **Plums:** 2 to 5 applications of 1½ to 2 quarts/acre from petal fall. Repeat applications at 10 to 14 day intervals. Water rate: 100 gallons/acre.
- **Potatoes:** To increase tuber size: A minimum of 2 applications of 2 quarts/acre during tuber bulking (as soon as first formed tuber are ½" in diameter) and following petiole analysis during tuber bulking. Allow 10 to 14 days between applications. Water rate: 20 gallons/acre.
- **Rice:** 2 quarts/acre applied at tillering and again at panicle initiation. Repeat 7 to 10 days later if necessary. Water rate: 3 to 40 gallons/acre.
- **Sorghum:** 2 quarts/acre at 4 to 8 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 3 to 20 gallons/acre.
- **Soybeans:** 1 to 2 quarts/acre when crop is 6 inches tall. Repeat if necessary at 10 to 14 day intervals. Water rate: 3 to 20 gallons/acre.
- **Spinach:** 2 quarts/acre at the 4 to 6 leaf stage. Water rate: 20 gallons/acre Squash (Field Grown): 2 quarts/acre at 4 leaf stage. Repeat if necessary 10 to 14 days later. Water rate: 50 gallons/acre.
- **Strawberries (Field Grown):** Non everbearing varieties: 3 applications of 2 quarts/acre from start of flowering. Repeat applications 7 to 10 day intervals. Everbearing varieties: 6 applications of 2 quarts/acre. Do not apply successive applications at intervals of less than 10 to 14 days. Water rate: 50 gallons/acre.
- **Sugar Beets:** 1 to 2 quarts/acre at the 4 to 6 leaf stage. For moderate to severe deficiency, repeat at 10 to 14 day intervals. Water rate: 20 gallons/acre.
- **Sweet Potato:** 2 quarts/acre one week after 100% emergence or transplanting. Repeat applications during tuber bulking at 10 to 14 day intervals. Also, apply at the same rate following recommendation from analysis. Water rate: 20 gallons/acre.
- **Tomatoes (Field Grown):** 2 quarts/acre. Apply at the 4 to 6 leaf stage with repeat applications at 10 to 14 day intervals if necessary. Water rate: 50 gallons/acre.
- **Water Melons (Field Grown):** 2 quarts/acre at the 4 to 6 leaf stage. Repeat at 10 to 14 day intervals. Water rate: 20 to 50 gallons/acre.
- **Zucchini (Field Grown):** 2 quarts/acre at the 4 to 6 leaf stage. Repeat at 10 to 14 day intervals. Water rate: 20 to 50 gallons/acre.

\*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.