




YaraAmplix™
ACTISIL®



Net contents:

**1.3 gallons
5 litres**

**Net weight
12.32 lb/
5.59 kg**

**Density
9.48 lb/gallon
1.12 kg/L**

Batch number:

NOTICE:

Before using this product, read the entire label.

ACTISIL®

A formulated product to strengthen crops.

Guaranteed Analysis:

Calcium (Ca) 2 %

Derived from Calcium Chloride.

ALSO CONTAINS NONPLANT FOOD INGREDIENT:

Soluble Silicon (Si) 0.5 %

Derived from choline-stabilized orthosilicic acid

Patent US 7968528

F1106

Yara North America, Inc

100 North Tampa Street, Suite 3800

Tampa, FL 33602 Customer service: 800-234-9376

Manufactured for Yara UK by Bio Minerals NV

Country of origin: Belgium

Danger

May be corrosive to metals. Causes severe skin burns and eye damage.

Do not breathe gas or vapour.

Wear protective gloves/clothing and eye/face protection.

Keep only in original container.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

IF ON SKIN (or HAIR): Take off immediately all contaminated clothing. Rinse skin with water.



YaraAmplix™ ACTISIL®

CROPS, APPLICATION RATES AND TIMINGS

FRUIT: (e.g. strawberries, apples, berries, kernel fruit and stone fruit)

Small Fruit and Berries (e.g. berries: raspberries):

Foliar Application: every week at the start of fruiting

- Number of treatments: 4-8

- Dose: 7 fl oz in 50 gallons/acre

Strawberries:

Foliar Application: every week at the start of flowering

- Number of treatments: 4

- Dose: 7 fl oz in 50 gallons/acre

Pome Fruit and Stone Fruit (e.g. apples, pears, cherries):

Foliar Application: every 7 - 10 days after flowering

- Number of treatments: 4-8

- Dose: 5 fl oz in 35 gallons/acre

VEGETABLES: (open field and under glass)

Fruiting vegetables: (e.g. tomatoes, peppers, eggplant, cucumbers, gherkins, zucchini, pumpkins)

Foliar Application: every week at the start of fruiting

- Number of treatments: 3-6

- Dose: 7 fl oz in 50 gallons/acre

Leafy vegetables: (e.g. lettuce, lamb's lettuce, endives, garden cress)

Foliar Application: every week upon development of leaf mass

- Number of treatments: 2-4

- Dose: 7 fl oz in 50 gallons/acre

Brassica vegetables: (e.g. cauliflower, broccoli, red cabbage)

Foliar Application: every week after planting

- Number of treatments: 2-4

- Dose: 5 fl oz in 35 gallons/acre

Foliar application.

Use 14 fl oz YaraAmplix™ ACTISIL® in 35 – 100 gallons water per acre for weekly application, as per requirements.

Fertigation application.

YaraAmplix™ ACTISIL® can also be applied via fertigation.

Use 1 pint YaraAmplix™ ACTISIL® in 250 gallons water.

STORAGE AND TRANSPORT

Store at room temperature in a sealed container protected from light.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

WARNING: Cancer and Reproductive Harm - www.P65Warnings.ca.gov

Only apply to the intended application site. Do not apply near water, storm drains, or drainage ditches or to any impervious surface. Do not apply if heavy rain is expected.

Choline-stabilized orthosilicic acid (ch-OSA®) is a product of Bio Minerals N.V. (Belgium). ch-OSA and Actisil are registered trademarks of Bio Minerals N.V.

